

CCA ADVANTAGE

*The Voice of the Certified Crop Adviser Program
American Society of Agronomy
www.agronomy.org/cca*



International Certified Crop Adviser Leadership for 2003-04

These are the newly elected International Certified Crop Adviser (CCA) chair, vice chair and regional representatives. Their profiles and comments on the important issues follow.

ICCA CHAIRMAN

Bob Beck, CCA, Pioneer Hi-Bred International's manager of technical training, Johnstown, IA, is the new chairman. His term of office begins in October and extends through September 2004.

His crop advising activities include developing fertilizer recommendations for crops grown in 15 states, and he also teaches courses for continuing education credits (CEUs). He is a certified professional soil scientist, a member of the American Society of Agronomy and a member of the Soil Science Society of America.

Beck says it is important that ICCA "really have the pulse of what the membership wants. That is so hard to do. Once you have a membership right at 15,000, you've done something right," he says.

Back in the late 1980s, he recalls, USDA believed that those with a "vested interest" in selling products for a



*Bob Beck, CCA
ICCA Chairman
Pioneer Hi-Bred*

company should not be allowed to advise growers. "We've had an impact on that; it's gone away for the most part," he says. CCAs "live, breathe and study" and bring credibility to the process of bringing the right information to their clients to make the right decisions, and that's a strength of the program, says Beck.

"We need to make sure we are indeed bringing programs that meet the needs of our customers. The weakness is whether we are fully cognizant of where we need to go to meet these needs. CCAs need to be aware and knowledgeable of controversial issues such as biotech crops, organic farming, etc., so you can bring options and best practices to suggest" to clients.

He would like to see the "T" in ICCA expanded during his tenure as chairman. Approaching a more international membership doesn't involve waiting for other countries to knock on ICCA's door. "We may have to make some real strong overtures to them. China is out there, as is the whole Asian theater. There's a tremendous interest," says Beck.

ICCA VICE CHAIRMAN

Stephen A. Dlugosz, CCA, is the new vice chairman of ICCA, with a term extending through September 2004. He is an agronomist for AgOne and Harvest Land Co-ops for east central Indiana and west central Ohio, with activities that include continuing education of sales and

support staff, in-field problem diagnosis, grower education and field days.

Dlugosz is working hard to add more value to the program to assist the individual CCA working in the field. "When I talk with co-workers and other CCAs about the program, I frequently hear their concerns that it is only a series of requirements consisting of an annual fee and continuing education units," says Dlugosz. The ICCA board is "very aware" of these concerns "and will be working hard to increase value not only for the individual CCA but for other participants in the food production system," he says. "Once a CCA recognizes how these individuals value the program, their personal value also will increase."

Another challenge is the consolidation within the various sectors of agriculture. "Good, devoted people in each state are what makes this program work. Finding these individuals to serve their local boards becomes more of a challenge every day," he says.

More emphasis on high-quality educational programs also interests him.



*Stephen A. Dlugosz
CCA, ICCA Vice
Chairman
AgOne and Harvest
Land Co-ops*


continued on page 16

The views expressed in this publication represent those of the authors. These views do not necessarily reflect endorsement by the American Society of Agronomy or any of the Society's programs. In addition, trade names are sometimes mentioned in this publication. No endorsement of these products or services by the American Society of Agronomy or any of its programs is intended, nor is any criticism implied of similar products or services not mentioned.

An electronic version of CCA Advantage, including the test, is available on our Web site: www.AgProfessional.com

for daily updates go to www.AgProfessional.com

AgProfessional • October 2003 • 15



continued from page 15

WESTERN REGION

Allen Smith, CCA, is the ICCA representative from the West. His term in office lasts three years. He is a crop consultant with Wilbur-Ellis Company, Quincy, WA, and consults on such crops as mint, small grains, forage crops, potatoes, carrots and various seed crops, focusing on fertility, moisture-related aspects of the crop and crop protection.



*Allen Smith, CCA
ICCA Board
Wilbur-Ellis*

The “key strength” of the CCA program is that it is more professional for growers, consumers and the general public, he says. And a CCA is more knowledgeable in fertility, “which is a must in today’s world.”

Smith believes his greatest challenge in the coming year will be to communicate with CCAs in the western region and get their inputs to pass on to the entire board.

“The role CCAs can play in helping resolve issues is to be informed and up to date,” he says. “We need to get the word out to the public that we as CCAs are using best management practices and will continue to do so.”

Smith has served on the CCA Columbia Basin Crop Consultants Association Board and Columbia Basin Crop Consultants Short Course Planning Committee and on the Washington State Pest Consultants Board.

NORTHEAST REGION

Terry Williams is the Northeast representative on the ICCA Board for a three-year term. He is a crop adviser for Royster-Clark Inc. in Dresher, PA.

“Developing and maintaining good lines of communication between crop advisers and the public seems to be our greatest challenge,” says Williams.

The greatest strength of the CCA Program, he believes, is “a group of

extremely knowledgeable and experienced professionals in the field of agriculture.” What CCA needs, he adds, is to focus on educating not only those involved in agriculture but the general public as well.

As a new board member, Williams hopes to broaden his knowledge as well as work with other board members to resolve issues facing agriculture. To do that, he plans to stay up to date on issues and technical advancements, attend educational seminars and suggest new seminars, speak with farmers and CCAs about the issues that concern them and bring these issues to the board.



*Terry Williams
CCA
ICCA Board
Royster-Clark*

Williams works with 25 growers, focusing on fertility, pest management and crop management.

NORTH CENTRAL REGION

Harold Watters, CCA, will be a North Central representative for two years. He is an extension agent, agricultural and natural resources/community development for the Ohio State University.

He says the CCA program “is at a fork in the road” and has been for some time. “Fewer and fewer candidates are taking exams and becoming CCAs,” he says.

Also, he says, “We need to consider the needs of the persons involved in their work and be much less concerned about getting a candidate through an exam.”

A lot of “very hard-working folks have given a lot to this program,” says Watters. “That history and the support of the American Society of Agronomy are the strengths of the organization.”



*Harold Watters
CCA
ICCA Board
Ohio State
University*

Getting CCAs approved as technical service providers (TSPs) “may have been the biggest boost for the program ever,” he says. “I want to see further efforts to help CCAs get involved as TSPs.”

CCAs have been and will continue to be educators and advisers, he adds. “They have the ability to observe, measure and evaluate the worth of a product, a practice or a concept. When something new comes along that may benefit the farmer, the CCA has the ability to measure that value.”

Waters is a member of the OSU Agronomic Crops Team, OSU Watershed Team and OSU Land Use Team.

Howard Brown is a North Central board representative for a term of three years. He is manager of agronomy services for Growmark and lives in Mahomet, IL.

Brown earned his bachelor’s degree from Southern Illinois University, a master’s from Purdue and a doctorate from the University of Illinois.

From 1980 to 1988 he worked in various positions for Corn Belt FS, Inc., Decatur, IL. He was field sales agronomist for Pioneer Hi-Bred International from 1988 to 1998, when he took his current position.

He makes in-field service calls for member cooperatives in Illinois, Iowa and Wisconsin, assists in assessing of problems and suggests corrective measures. He also works with in-field CCA training and works closely with USDA’s Natural Resources Conservation Service in Illinois, Iowa and Wisconsin to stay current with issues on TSP.

He was chair of the CCA Exam Committee and the Illinois CCA Board, and is a CCA Board ex-officio member and a member of the Illinois Chemical and Fertilizer Association agronomy committee.



*Dr. Howard
Brown, CCA
ICCA Board
Growmark*



Chairman's Corner

Think Global, Act Local

By Tom Bruulsema, Chair, International CCA Board

It's hard to believe that my year as chairman of the ICCA Board is already coming to a close. As I write this, I'm still looking forward to a couple of exciting challenges this fall. It has indeed been a pleasure working with the fine people you've elected. I've highly appreciated the advice and continued involvement of Pat McConnell, past-chair, and I'm confident your leadership is in good hands with incoming chair Bob Beck and, following him, Steve Dlugosz.

I'd like to spend some time reflecting on the value of the international dimension of the CCA program. Why have an International Board in the first place?

federal involvement. Nutrient management plans, conservation plans and other environmental protections are evolving to meet both local and federal standards, in both the U.S. and Canada. The farm bill in the U.S. and the Agricultural Policy Framework in Canada will increasingly impinge on these issues.

The food on our ultimate customer's plate comes from across the continent. And that customer needs assurance that the product — food, fiber or fuel — is safe, has quality and was produced in an environmentally friendly manner. We won't get the chance to explain the detailed efforts expended to achieve those production standards. But if we can make



Tom Bruulsema, Ph.D.
Chair, ICCA Board

What's the value in holding a designation the same as the one someone else holds a couple thousand miles away, advising producers of crops you've never seen? In today's world, the value is enormous!

Our programs are founded on enormous efforts expended by people who have top expertise at the local level — be they from university, industry or government agency. The local boards have been a strong part of the program's success. So much so, it's possible to question the value of the international component. I've been there! But there are plenty of reasons why it's essential to have a strong International Board, and a well-standardized program.

Our major issues have always had both state/provincial and federal jurisdictions — and the trend is toward stronger

our certification a recognized part of a system of standardized, branded agricultural production, our credentials will rise in value and contribute to the assurance. We need to dream of a time ahead when the involvement of a Certified Crop Adviser forms part of the requirement for a recognized brand of product and standard of quality — and work to make that dream a reality.

What are the benefits of a strong International program?

- A one-stop shop where anyone can find a CCA.

- A consistent message on what CCA is all about.
- Expertise in program development and management.
- A strong standard of quality in the exams.
- Ability to contribute toward meeting international trade standards.
- Mobility — your certification moves with you when you take advantage of new opportunities.
- Learning about crop production in other areas — even if you've never seen them before — helps you better your understanding of the principles of agronomy.
- Efficiency in managing your program.

The inputs we deliver also come from across the continent, and indeed from beyond its borders. From a product stewardship perspective, we need consistent standards, whether they are applied in Alaska, Newfoundland, Florida or California.

Let's continue a strong partnership between local and international leadership of this truly successful program.

Tom Bruulsema is Director, Eastern Canada and Northeast U.S. region with the Potash & Phosphate Institute. He lives in Guelph, Ontario, Canada.



Continuing Education Self-Study Course

Soil & Water Management



Relationship of Soil Test Phosphorus and Sampling Depth to Runoff Phosphorus in Calcareous and Noncalcareous Soils

Earn one CEU!

All CCAs may earn up to 20 Continuing Education Units (CEUs) per two-year cycle as board-approved self-study articles which will include CCA Advantage articles. The CCA CEU logo (above) marks all pre-approved material, with the CEU value indicated by the number in the middle. To receive one CEU in soil and water management, read this article, fill out the attached exam and mail the tear-out form, along with \$10, to the American Society of Agronomy.

By H. A. Torbert, T. C. Daniel, J. L. Lemunyon, and R. M. Jones

In a 1998 U.S. EPA report, agricultural nonpoint source pollution was estimated to cause 60%, 50% and 34% of river, lake and estuarine impairment, respectively. The greatest potential for nonpoint P contribution to surface waters usually occurs in watersheds with intensive animal production. Manure collected from concentrated animal feeding operations (CAFO) has traditionally been applied to fields near the operation. However, long-term manure application to soils at rates in excess of crop uptake can result in elevated soil P levels.

Research in 1993 showed that the contribution of P from soils with elevated levels of soil test P is potentially more important and difficult to manage than improper land application of animal manure. This study found that elevated soil test P levels were responsible for 65% to 90% of annual P loss from the watershed even when a major surface

runoff event occurred one day after manure was applied to fescue pasture.

Recent EPA draft guidelines for manure applications for CAFOs limit application to “threshold P holding capacity of all major soil types within the land application areas.” However, the threshold P levels of soils are yet to be developed. Recent research has shown that the level of soil P is directly related to runoff losses of P, which would indicate that a threshold level could be developed, but studies have demonstrated that the relationship between the level of soil P and runoff P varies markedly depending on the soil type. Further, it cannot be assumed that soil tests designed for crop production can be used to predict surface runoff enrichment potential.

Using the relationship of soil test P to runoff losses of P could be a valuable management tool for protecting watersheds from excessive nutrient loading. However, the potential of these relationships has not been fully developed, especially for calcareous soils. The objective of this project was to examine the relationships between soil test P levels and runoff losses of P for benchmark soils in the Bosque River watershed in Texas and to examine the practical aspects of the data (i.e., depth of sampling and P extractant used) for potential use in development of a soil test for environmental P losses.

Materials and Methods

Rainfall simulations were conducted on four soils representative of the major soil types in the Bosque River basin in Texas. The soils included two noncalcareous soils, a Windthorst sandy loam and a Blanket clay loam, and two calcareous

soils, a Purves clay and a Houston Black clay. The soils were under permanent Bermuda grass pasture.


At each site, six surface runoff plots were constructed and a collection gutter was placed at the downslope edge to divert runoff to a pit where samples could be collected. A range in the level of soil test P was established by surface application of dairy manure from a local dairy operation. A target range of soil test P levels of 0, 60, 120, 180, 240 and 360 mg kg⁻¹ was used. Following application of manure over an 18-month period, approximately six months were allowed before initiation of rainfall simulation so the effect of surface manure application on runoff P concentration would be minimized. The grass was mowed periodically.

A composite soil sample was collected from each plot immediately after simulated rain applications to permit correlation of soil test and surface runoff P levels. Soil cores were taken at 0- to 2.5-, 0- to 5- and 0- to 15-cm depths.

A rain simulator generated 30 minutes of surface runoff from each plot. Runoff samples were filtered, acidified to pH 2 with HCl and frozen until analyzed for dissolved molybdate reactive phosphorus (DMRP). Rainfall simulations were repeated three times at each site so that regression analysis could be used to evaluate the relationship between soil test P and surface runoff P.

Results

The soils were very different with respect to texture and CaCO₃ content, especially in the upper few centimeters, which is expected to have the greatest influence on runoff P concentrations. While CaCO₃ is present in all four soils, levels in the sur-



face horizon of the Houston Black and Purves soils (considered calcareous) are 10-fold higher than the Windthorst and the Blanket soils.

Soil samples collected at the four locations indicated that the concentrations of extractable P were reasonably close to the target levels of 0, 60, 120, 180, 240 and 360 mg kg⁻¹.

Analysis of runoff P concentrations across all four soil types ranged from 0.023 to 1.73 mg L⁻¹ for DMRP and from 0.044 to 1.8 mg L⁻¹ for total P. For all soils, runoff P concentrations increased as the level of soil test P increased.

However, the amount of particulate P (particulate P = total P – DMRP) in runoff remained relatively constant with increasing levels of P in runoff and contributed a very small portion of the total P in runoff at the higher levels of runoff P observed. Since the greatest potential for increased eutrophication of surface water results from DMRP components, regression analysis presented here will be for DMRP.

It has been speculated that a “change point” may exist for soil test P concentrations where increasing P application (i.e., manure application) contributes increasing concentrations of DMRP to runoff. Within the range of soil test P used in this study, no breakpoints for soil test P concentrations were observed.

Soil Depth. A strong relationship exists between soil test P and DMRP in surface runoff with both soil test extractions using water and Mehlich III. Similar results have been reported by others, and our results are further evidence that an environmental soil test can be developed that relates the level of P measured in soil to the amount susceptible to losses in surface runoff.

While similar, significant regression equations were observed for all four soils and all three soil depths, examination of the differences and similarities can provide some indications as to the best soil test for P relating to potential environmental loss of P. For example, while the relative response was different between the four soil types, the relative response between the four soil types for changing soil depth was consistent. The greater the soil depth, the steeper the curve response to increasing soil test P. We believe this is

primarily a factor of dilution of the manure P spread on the surface of the established pasture with the inclusion of deeper sampling increments. In this system, the manure P must move through the sod and down through the soil profile. Consequently, the highest concentrations of P were observed in the soil surface and decreased with increasing depth.

The Windthorst soil is a sandy loam soil compared with the other three, which were clay and clay loams. With the sandy soil, manure additions to the soil surface can move through the soil profile more quickly; consequently, there is less dilution effect of the soil test P values. This resulted in a reduction of the relative change in steepness of regression equations for soil depth in the Windthorst soil compared with the heavier-textured soil types.

While not always the case, the best fit for the regression lines for both Mehlich III and water-extractable P was generally observed with soil collected from the 0- to 5-cm depth. It is believed that this was caused by competing phenomena relative to manure application on the soil surface.


Statistical techniques were used to compare differences between regression equations at each soil depth between the four soil types. With the water-extractable P, significant differences were observed between the regression equations of the four soil types for both the 0- to 2.5- and 0- to 5-cm depths but not the 0- to 15-cm depth. At the 0- to 15-cm depth, only the soil P concentration level was found to have a significant effect on the regression equations. This indicates that the equations for each soil type would be parallel with a different intercept for each soil type. However, given the response observed at the other two soil depths, this response would not seem to be realistic. It is believed that this response at the 0- to 15-cm depth is actually a diminishment in the effectiveness to describe the relationship, with only the strongest component (P concentration) remaining significant compared with the responses observed at the 0- to 2.5- and 0- to 5-cm depths. At the 0- to 15-cm depth, while the clear relationship between soil test P level and runoff P losses could be observed, differences between the soil types were not significant. While this

may be desirable for providing a “one fits all” line for making an environmental P soil test, it would potentially eliminate important differences in soils for management purposes.

At the 0- to 2.5-cm depth, a significant difference was observed for water-extractable P concentration and soil type, but only the Houston Black soil was significantly different from the other three. This lack of significance for the other soil types was probably due to increased variability in the P concentration data at the 0- to 2.5-cm depth. At the 0- to 5-cm depth, the Blanket, Houston Black and Purves soils were all significantly different from each other. The Windthorst soil was significantly different from the Houston Black soil, but was only close to being significantly different from the other two soil types.

Similar results were observed for differences among the regression equations developed for the Mehlich III extractant. As with the water-extractable P at the 0- to 15-cm depth, P concentration was the only significant effect observed. However, with the Mehlich III extractant, differences could be observed between soil types at the 0- to 2.5- and 0- to 5-cm depths. This indicates that Mehlich III may be less variable with P concentration data at the 0- to 2.5-cm depth for developing these types of relationships compared with water-soluble P extraction. However, both extractants resulted in a highly significant relationship at the 0- to 5-cm depth, indicating that this may be the best depth for examining DMRP losses regardless of the extractant used. As was seen with the water-extractable P, at the 0- to 5-cm depth, the Blanket, Houston Black and Purves soils were significantly different and the Windthorst soil was close to being significantly different from the Purves and Blanket soil types.

Calcareous versus Noncalcareous Soil. Comparison of the four sites indicates another difference among the regression equations. The two calcareous soils have a much lower concentration of DMRP in runoff at all levels of soil test P compared with the two noncalcareous soils. With the noncalcareous soils, DMRP concentration in runoff varies from 0.2 to 1.5 mg L⁻¹, while the DMRP



concentration in the calcareous soils started well below 0.2 mg L^{-1} and at the highest soil test P level only reached a concentration of 0.6 mg L^{-1} . In fact, the Houston Black soil only reached a value of 0.4 mg L^{-1} at the highest soil test P concentration. These results have important implications for regulations using soil test P values because these relatively large differences in DMRP losses were observed at the same measurement of soil test P using the same laboratory techniques for determination.

Calcareous soils by definition contain sufficient free CaCO_3 to effervesce visibly when treated with cold 0.1 M HCl . Soluble P will react with Ca in soil to form insoluble minerals such as hydroxyapatite and fluorapatite. Therefore, calcareous soils will probably contain less soluble P at higher levels of total soil P (higher manure application). The presence of free CaCO_3 in calcareous soils will probably reduce the amount of soluble P present in soil and prevent it from being released into runoff.

Soil P extractants were developed to measure the amount of P that would be released during the growing season for plant production. Therefore, most extractants will dissolve some of the insoluble Ca-phosphate minerals present in soil. The data presented here demonstrate that while the P concentrations present in extractants are directly related to the potential P release into runoff, the relationship can be highly influenced by the presence of soil components that form insoluble P minerals.

While measuring differences between soil types is important, it is necessary to develop a practical soil test P for environmental concerns. Therefore, predictive equations are needed that will work satisfactorily for groups of soils to predict DMRP in runoff. In this study, an attempt was made to develop regression equations by grouping the two calcareous soils and the two noncalcareous soils. Significant regression lines were developed for comparing soils for both the water-extractable and the Mehlich III extractants, but not for all three soil depths. With the Mehlich III data, no significant regression lines could be developed for the 0- to 2.5-cm depth, and the regression equations developed at the 0-

to 15-cm depth were not significantly different from each other. However, at the 0- to 5-cm depth, regression equations were developed with reasonable r^2 values and a highly significant soil type effect. With the water-extractable P, regression lines could be developed at all depths, but no significant differences were observed with the 0- to 15-cm depth. A significant difference was observed between the regression models for the 0- to 2.5 depth, but only at the $\alpha = 0.1$ probability level ($P = 0.097$). A highly significant difference was observed between the two models at the 0- to 5-cm depth ($P = 0.004$) with reasonable r^2 values for the two lines.

Discussion

The data presented in this study indicate significant differences in the potential for runoff losses of DMRP at the same concentration of soil test P between different soil types within the same watershed. Also, the data indicate there is a potential to group classes of soil types for their soil DMRP loss potential. These differences could be used in tools such as the Phosphorus Index for manure management.

If a soil test for P is to be used as a management tool for land application of manure, soils will need to be grouped into reasonable management categories and reliable predictive equations for potential P loss developed for those soil categories. Grouping soils by chemical characteristics (calcareous vs. noncalcareous) allowed development of significant predictive equations.

Differences observed between depths of soil sampling may also be very important in developing a P soil test for environmental response purposes. Besides the physical problem of obtaining a consistent soil sample on the soil surface, soil sampling at the 0- to 2.5-cm depth tended to increase variability, indicating that sampling at this depth may be problematic.


Likewise, soil sampling at the 0- to 15-cm depth may also be problematic for soil testing for potential environmental P losses. The potential problem with the data developed for the 0- to 15-cm depth is the steepness of the curve, with differences between agronomic response levels and excessive levels being small. The

result of a steep response curve is that small differences in soil test P concentration could result in huge differences in interpretation. This would in turn put more pressure on sampling methods in the field and measuring equipment in the laboratory to assure accuracy of measurements. Since large changes in the field would be measured with small differences in soil test P results, it would be difficult for operators to follow the progress of P enrichment in their fields. In addition, a reduction in the capacity to distinguish difference between soil types may result in difficulty for developing predictive equations over a large grouping of soil types.

The data indicate that sampling at the 0- to 5-cm depth may arrive at the best results both from limiting management problems and by increasing the likelihood of developing reliable predictive equations across a large number of soil types.

The data presented in this study demonstrate two important points: (1) differences exist in soil that may affect the level of DMRP contributed to the environment at the same level of soil test P and (2) these differences could potentially be used in the management of CAFOs to reduce DMRP levels in runoff. Also, the data demonstrate that there may be chemical or physical characteristics of soils that can be used to develop application criteria in predictive tools such as the Phosphorus Index.

Editor's note: Content was adapted from the paper "Relationship of Soil Test Phosphorus and Sampling Depth to Runoff Phosphorus in Calcareous and Noncalcareous Soils," which was published in the Journal Environ. Qual, 2002 31 and is courtesy of the authors H. A. Torbert, T. C. Daniel, J. L. Lemunyon, and R. M. Jones.



Get a CEU!

This exam is worth 1 CEU in **Soil and Water Management**. An exam score of 70% or higher will earn CEU credit. The International CCA program has approved self-study CEUs for 20 of the 40 CEUs required in the two-year cycle.

DIRECTIONS

1. Read the self-study article on pages 18-20 carefully.
2. Answer the questions by clearly marking an "X" in the box next to the best answer for each question.
3. Complete the self-study exam registration form on the back of this page.
4. Clip out this self-study examination page, fold and place in envelope.
5. Enclose a check for \$10.00 made payable to the American Society of Agronomy, for processing fees. Payment in U.S. funds only.
6. **Mail your self-study exam and fee to:**
ASA c/o CCA Self-Study Exam, 677 S. Segoe Road, Madison, WI 53711. *Please allow 60 days for processing.*
7. An electronic version of this test is also available at www.AgProfessional.com. Go to the Certified Crop Advisers section (lefthand column) and access the "CCA Advantage" link.

Relationship of Soil Test Phosphorus and Sampling Depth to Runoff Phosphorus in Calcareous and Noncalcareous Soils

October Self-Study Examination

1. **Impairment of rivers, lakes and estuaries by agricultural nonpoint source pollution has been reported at:**
 - a. 60%, 50% and 34% respectively.
 - b. 55%, 45% and 32% respectively.
 - c. 50%, 40% and 30% respectively.
 - d. 45%, 35% and 28% respectively.
2. **Usually the greatest potential for nonpoint P contribution to surface waters occurs in watersheds with:**
 - a. intensive urban development.
 - b. intensive crop production.
 - c. intensive animal production.
 - d. extensive wildlife habitat.
3. **Elevated soil P levels can be the result of:**
 - a. poor soil structure.
 - b. long-term manure application to soils at rates in excess of crop uptake.
 - c. long-term manure application to soils at a rate equivalent to crop uptake.
 - d. long-term fallow.
4. **Studies have found that the relationship between the level of soil P and runoff P varies depending on:**
 - a. soil type.
 - b. sample techniques.
 - c. statistical analysis.
 - d. topography.
5. **Runoff samples were analyzed for:**
 - a. runoff P.
 - b. particulate P.
 - c. total P.
 - d. dissolved molybdate reactive phosphorus.
6. **The greatest potential for increased eutrophication of surface waters results from:**
 - a. total P.
 - b. particulate P.
 - c. DMRP components.
 - d. runoff P.
7. **The study found the best depth for examining DMRP losses regardless of the extractant used was:**
 - a. 0 to 1 cm.
 - b. 0 to 2.5 cm.
 - c. 0 to 5 cm.
 - d. 0 to 15 cm.
8. **Compared to noncalcareous soil, the two calcareous soils have a much lower concentration of DMRP in runoff at:**
 - a. the 0 to 2.5 cm zone.
 - b. the 0 to 5 cm zone.
 - c. the 0 to 15 cm zone.
 - d. all depths.

Over

DETACH HERE

Continuing Education Self-Study Test

Soil and Water Management Test (continued)




9. A regression equation with a reasonable r^2 was developed for:

- a. the 0 to 2.5 cm depth.
- b. the 0 to 5 cm depth.
- c. the 0 to 15 cm depth.
- d. all of the sampling depths used in the study.

10. The data in this study demonstrate that significant predictive equations could be developed:

- a. using the 0 to 2.5 cm sampling depth.
- b. by grouping soils by chemical characteristics.
- c. by ignoring runoff losses of DMRP.
- d. by grouping soils according to geography.



SELF-STUDY EXAM REGISTRATION FORM

Name: _____
Address: _____
City: _____ State/Province: _____ Zip: _____
CCA Certification #: _____
Credit Card #: _____ Type of Card: Visa Mastercard Discovery Am Express
Expiration Date _____ Name on Card: _____

A \$2 processing fee will be added to all credit card charges, or enclose \$10 check payable to American Society of Agronomy.
X

Signature of Registrant as it appears on Code of Ethics

I certify that I alone completed this self-study course and recognize that an ethics violation may revoke my CCA status.

This exam issued October 2003 expires October 2006.

SELF-STUDY EXAM EVALUATION FORM

Rating Scale: 1=Poor 5=Excellent

Information presented will be useful in my daily crop advising activities: 1 2 3 4 5

Information was organized and logical: 1 2 3 4 5

Graphics/tables were appropriate and enhanced my learning: 1 2 3 4 5

I was stimulated to think how to use and apply the information presented: 1 2 3 4 5

This article addressed the stated competency area and performance objective(s): 1 2 3 4 5

Briefly explain any "1" ratings: _____

Topics you would like to see addressed in future self-study materials: _____

DETACH HERE